#### 2-DAY GOLYTELY SPLIT PREP

# Follow these instructions to prep for your procedure. Tulsa VA Endoscopy Clinic M-F 0700-1530 (918) 252-8000 ext 29042 or 29043 24/7 Colonoscopy Instruction Line at 918 577 4900

If you have had cardiac stents placed within the last 12 months please call our clinic.

### **7 DAYS BEFORE YOUR PROCEDURE:**

**STOP** taking **blood thinners:** EFFIENT (Prasugrel), Brilinta (ticagrelor), and/or PLAVIX (Clopidogrel).

**STOP** weight loss meds: (such as OZEMPIC, WEGOVY, MOUNJARO, and PHENTERMINE).

## **5 DAYS BEFORE YOUR PROCEDURE:**

**STOP** taking **blood thinners:** WARFARIN (Coumadin), <u>Aspirin</u>, any NSAIDS, such as: Ibuprofen, Advil, Naproxen, Meloxicam, Diclofenac.

STOP taking FISH OIL, FIBER, IRON NUTS, AND SEEDS.

# 2 DAYS BEFORE YOUR PROCEDURE:

**STOP** taking **blood thinners:** Pradaxa (dabigatran); Xarelto (rivaroxaban); Eliquis (apixaban).

# **FOLLOW THESE ENDOSCOPY INSTRUCTIONS:**

(not the manufacturer instructions on the colon prep bottle)

## BEGIN 2 DAYS of CLEAR LIQUIDS ONLY, NO SOLID FOOD FOR 2 DAYS.

Clear liquids are items that you can see-through, such as: water, tea, black coffee, broth, apple juice, Gatorade, jello, popsicles.

**NO** red or orange colored items.

NO dairy products

NO alcohol

NO marijuana.

**1PM**: Mix up 1st gallon of GoLytely prep with water only and place in the refrigerator.

**4PM**: Begin drinking the GoLytely - 8 oz glass every 15 minutes until gone (**FINISH ENTIRE GALLON** even if stools are clear). If you become nauseous, wait 30-60 minutes then resume prep.

You may take your medications as prescribed, **EXCEPT** for those mentioned above.

\*\*\*\*FLIP THE PAGE TO CONTINUE WITH ADDITIONAL PREP INSTRUCTIONS

YOU AND YOUR LICENSED DRIVER MUST CHECK-IN TOGETHER. THE DRIVER MUST STAY AT THE CLINIC UNTIL YOU ARE DISCHARGED. THEY CANNOT DROP YOU OFF AND LEAVE OR YOU WILL BE RESCHEDULED.

#### 2-DAY GOLYTELY SPLIT PREP

# Follow these instructions to prep for your procedure. Tulsa VA Endoscopy Clinic M-F 0700-1530 (918) 252-8000 ext 29042 or 29043 24/7 Colonoscopy Instruction Line at 918 577 4900

## **1 DAY BEFORE YOUR PROCEDURE:**

Clear liquids ONLY all day again! Stay hydrated.

**1PM**: Mix up 2nd gallon of GoLytely prep with water and put in the refrigerator.

**4PM**: Drink **only half** the gallon of Golytely - 8 oz glass every 15 minutes; until half of the gallon is completed. If you feel nauseous, wait 30-60 minutes then resume drinking your prep. Place remaining half gallon in the refrigerator (to be completed the morning of your procedure).

You may take your medications as prescribed, **EXCEPT** for those mentioned above.

#### IF YOU ARE A DIABETIC:

HOLD EVENING DOSE OF ORAL DIABETIC MED(S)

TAKE ONLY 1/2 YOUR USUAL EVENING DOSE OF LONG-ACTING INSULIN.

DO NOT TAKE ANY SHORT ACTING INSULIN THE DAY BEFORE PROCEDURE OR THE MORNING OF.

Monitor your blood sugar to maintain a reading of 100 or higher. Drink apple juice, Gatorade (not red or orange) 7-Up, or Sprite to help maintain your blood sugar levels greater than 100.

**DO NOT** have ANYTHING by mouth after **MIDNIGHT**, **EXCEPT ONLY your prep as directed**; including **all** oral tobacco products, mints, chewing gum, etc.

#### THE MORNING OF YOUR PROCEDURE:

If you take Blood Pressure meds and/or Seizure meds, take with ONLY small amount of water.

**DO NOT** take any DIABETIC medications and **NO** insulin.

**DO NOT** have anything else by mouth except for the prep as directed.

DO NOT use oral tobacco products, chewing gum, marijuana, gummies, mints, coffee, etc.

## **4 HOURS BEFORE YOUR APPOINTMENT:**

Drink **ONLY** the remaining 1/2 gallon of Golytely - 8 oz glass every 15 minutes. **This must be <u>completed</u> at least <u>2hrs before</u> your appointment time. Then, <b>NOTHING** by mouth for a **total of <u>2</u>**hrs before your appointment! **FINISH ENTIRE GALLON** even if stools are clear.

(Example – for a 1000 procedure time, prep **must be** <u>COMPLETED</u> by 0800, giving you a 2 hour window in which you have **nothing by mouth**. Allow yourself enough time to complete it as directed.)

By splitting the prep in this way increases your chance of cleaning out the right side of your colon timely and having an excellent exam; decreasing the chance of missing a significant finding or having to redo the procedure due to poor pep quality.

YOU AND YOUR LICENSED DRIVER MUST CHECK-IN TOGETHER. THE DRIVER MUST STAY AT THE CLINIC UNTIL YOU ARE DISCHARGED. THEY CANNOT DROP YOU OFF AND LEAVE OR YOU WILL BE RESCHEDULED.